

VORONOI SUMMATION FORMULAE ON $GL(n)$

Fan Zhou

February 4, 2015

Abstract

We discover new Voronoi formulae for automorphic forms on $GL(n)$ for $n \geq 4$. There are $[n/2]$ different Voronoi formulae on $GL(n)$, which are Poisson summation formulae weighted by Fourier coefficients of the automorphic form with twists by some hyper-Kloosterman sums.

MSC: 11F30 (Primary), 11F55 (Secondary)

1 Introduction

A Voronoi summation formula of an automorphic form is a Poisson summation formula weighted by Fourier coefficients of the automorphic form with twists by some additive arithmetic functions. The Voronoi formula of an automorphic form on $GL(2)$ is well known and has a lot of application.

A Voronoi formula of an automorphic form on $GL(3)$ was firstly proved by Miller and Schmid in [MiSc1]. Later, a Voronoi formula was established for $GL(n)$ for $n \geq 4$ in [MiSc2], [GoLi1], and [GoLi2]. An adelic version was established in [IcTe]. The Voronoi formula has important application, such as a nontrivial bound on the exponential sum in [Mi] and subconvexity bounds in [Li].

More recently, Li and Miller established a different type of Voronoi formula on $GL(4)$ ¹. Their formula is an equality between a weighted sum of Fourier coefficients of an automorphic form on $GL(4)$ and a dual sum, both twisted by classical Kloosterman sums. This formula is different from those in [MiSc2], [GoLi1], [GoLi2], and [IcTe] and inspires us to develop this paper.

In this paper, we discover more Voronoi formulae on $GL(n)$ when $n \geq 4$. One of these new formulae overlaps the formula of Li and Miller. Our formula on $GL(n)$ is an equality between a sum of Fourier coefficients of an automorphic form twisted by hyper-Kloosterman sums of dimension $(k-1)$ and a dual sum twisted by hyper-Kloosterman sums of dimension $(n-k-1)$. The Voronoi formula of [MiSc1], [MiSc2], [GoLi1], and [GoLi2] are the case of $k=1$. Li and Miller's Voronoi formula on $GL(4)$ is the case of $n=4$ and $k=2$. For an automorphic form on $GL(n)$, there are $[n/2]$ different Voronoi formulae, which correspond to $k=1, \dots, [n/2]$ respectively.

¹Their result was announced at *Representation Theory, Automorphic Forms, and Complex Geometry, a conference in honor of the 70th birthday of Wilfried Schmid* in May 2013. The slides of Li's talk are available in the link: http://www.math.harvard.edu/conferences/schmid_2013/docs/li.pdf

For an integer $k \geq 1$ define the $(k-1)$ -dimensional hyper-Kloosterman sum by

$$Kl_k(m, q) = \sum_{\substack{x_1, \dots, x_{k-1} \pmod q \\ (x_1, q) = \dots = (x_{k-1}, q) = 1}} e \left(\frac{x_1 + \dots + x_{k-1} + m \overline{x_1 \dots x_{k-1}}}{q} \right)$$

for a prime number q . We state our formulae in the following.

Theorem 1.1. *Let π be an even Maass cusp form for $SL(n, \mathbb{Z})$ of the spectral parameter $(\lambda_1, \dots, \lambda_n) \in \mathbb{C}^n$ and $n \geq 2$. Let $A(m_1, \dots, m_{n-1})$ be the Fourier coefficient of π . Let $\omega \in C_c^\infty(0, \infty)$ be a test function with $\tilde{\omega}$ its Mellin transform and let q be a prime number, a and $\bar{a}$ two integers with $a\bar{a} \equiv 1 \pmod q$. For $1 \leq k \leq n-1$, we have*

$$\begin{aligned} & \sum_{m=1}^{\infty} A(1, \dots, 1, m) Kl_k(am, q) \omega(m) \\ & + \sum_{2 \leq l \leq k} \sum_{m=1}^{\infty} (-1)^{l+k} q^{l-1} A(1, \dots, 1, \overbrace{q, 1, \dots, 1}^{\text{position } l}, m) \omega(mq^l) \\ & = \frac{q^k}{2} \sum_{m=1}^{\infty} \frac{A(m, 1, \dots, 1)}{m} \\ & \times \left(Kl_{n-k}(\bar{a}m, q) (\Omega_+ + \Omega_-) \left(\frac{m}{q^n} \right) + Kl_{n-k}(-\bar{a}m, q) (\Omega_+ - \Omega_-) \left(\frac{m}{q^n} \right) \right) \\ & + \sum_{2 \leq l \leq n-k} \sum_{m=1}^{\infty} (-1)^{n-k+l} q^{k-1} \frac{A(\overbrace{m, 1, \dots, 1, q}^{\text{position } l}, 1, \dots, 1)}{m} \Omega_+ \left(\frac{m}{q^{n-l}} \right), \end{aligned}$$

where $\Omega_{\pm}$ is defined as

$$\Omega_+(x) = \frac{1}{2\pi i} \int_{\Re s = -\sigma} \tilde{\omega}(s) \pi^{-n(1/2-s)} \prod_{j=1}^n \Gamma \left(\frac{1-s-\bar{\lambda}_j}{2} \right) \Gamma \left(\frac{s-\lambda_j}{2} \right)^{-1} x^s ds$$

and

$$\Omega_-(x) = \frac{1}{2\pi i} \int_{\Re s = -\sigma} \tilde{\omega}(s) i^{-n} \pi^{-n(1/2-s)} \prod_{j=1}^n \Gamma \left(\frac{2-s-\bar{\lambda}_j}{2} \right) \Gamma \left(\frac{s+1-\lambda_j}{2} \right)^{-1} x^s ds.$$

Proof. This is literally the sum of Theorem 3.5 and Theorem 4.2. $\square$

Our proof is based on the functional equations of the twisted L -functions of π by (multiplicative) Dirichlet characters. The relationship between the Voronoi formulae and the functional equations of L -functions is known to experts. The same method was used in Section 4 of [GoLi1] and in proving Lemma 1.3 in [BuKh] for a special case on $GL(3)$.

Our proof does not depend on automorphicity but depends only on the family of the functional equations of $GL(1)$ -twisted L -functions. Functional equations of $GL(1)$ twists are consequences of automorphicity but they do not imply automorphicity on

$GL(n)$ when $n \geq 3$. In view of the converse theorem on $GL(n)$ of Cogdell and Piatetski-Shapiro [CoPS], a proof based on $GL(1)$ -twisted functional equations is stronger than a proof based on automorphy.

Our theorem and proof can be easily applied to many conjectural functorial lifts. For example, although the Rankin-Selberg convolutions on $GL(m) \times GL(n)$ are not generally known to be automorphic on $GL(m \times n)$, except the few cases of $GL(2) \times GL(2)$ and $GL(2) \times GL(3)$, the functional equations of $GL(m) \times GL(n) \times GL(1)$ are well known. Thus, we have the Voronoi formulae for the Rankin-Selberg convolutions on $GL(m) \times GL(n)$. Considering the Langlands-Shahidi method and the general Rankin-Selberg method, there are numerous functorial cases to have Voronoi formulae without being known to be automorphic.

The obvious weakness of our result is that the denominator q in the hyper-Kloosterman sums has to be a prime number. More general formulae with arbitrary integer q awaits exploration.

Acknowledgment

The author wants to thank Wenzhi Luo, Roman Holowinsky, and Eyal Kaplan for helpful discussion. The author wants to thank Dorian Goldfeld for introducing this topic to him.

2 Background

Let π be an even Maass cusp form for $SL(n, \mathbb{Z})$ with the spectral parameter $(\lambda_1, \dots, \lambda_n) \in \mathbb{C}^n$. Let $A(*, \dots, *)$ be the Fourier-Whittaker coefficient of π . We have

$$A(\pm m_1, \dots, \pm m_{n-1}) = A(m_1, \dots, m_{n-1}),$$

$$\overline{A(m_1, \dots, m_{n-1})} = A(m_{n-1}, \dots, m_1),$$

and $A(1, \dots, 1) = 1$. We refer to [Go] for the definitions and the basic results of Maass forms for $SL(n, \mathbb{Z})$. The dual (contragredient) Maass form of π is denoted by $\tilde{\pi}$. Let $B(*, \dots, *)$ be the Fourier-Whittaker coefficient of $\tilde{\pi}$ and we have

$$\overline{A(m_1, \dots, m_{n-1})} = B(m_{n-1}, \dots, m_1).$$

The standard L -function of π is

$$L(s, \pi) = \sum_{m=1}^{\infty} \frac{A(1, \dots, 1, m)}{m^s}.$$

It satisfies the functional equation

$$L(s, \pi) = G_+(s) L(1-s, \tilde{\pi}), \tag{1}$$

where

$$G_+(s) = \pi^{-n(1/2-s)} \prod_{j=1}^n \Gamma\left(\frac{1-s-\bar{\lambda}_j}{2}\right) \Gamma\left(\frac{s-\lambda_j}{2}\right)^{-1}.$$

Let q be a prime number. Let ψ be a nontrivial Dirichlet character mod q and $\tau(\psi)$ its Gauss sum. The ψ -twisted L -function is

$$L(s, \pi \times \psi) = \sum_{m=1}^{\infty} \frac{\psi(m) A(1, \dots, 1, m)}{m^s}.$$

Let k be an integer with $1 \leq k \leq n-1$. In the case of $\psi(-1) = 1$, the twisted L -function satisfies the functional equation

$$L(s, \pi \times \psi) = \tau(\psi)^n q^{-ns} G_+(s) L(1-s, \tilde{\pi} \times \bar{\psi}),$$

which can be rewritten as

$$\tau(\bar{\psi})^k L(s, \pi \times \psi) = \tau(\psi)^{n-k} q^{k-n} G_+(s) L(1-s, \tilde{\pi} \times \bar{\psi}). \quad (2)$$

In the case of $\psi(-1) = -1$, the twisted L -function satisfies the functional equation

$$L(s, \pi \times \psi) = \tau(\psi)^n q^{-ns} G_-(s) L(1-s, \tilde{\pi} \times \bar{\psi}),$$

which can be rewritten as

$$\tau(\bar{\psi})^k L(s, \pi \times \psi) = \tau(\psi)^{n-k} q^{k-n} G_-(s) L(1-s, \tilde{\pi} \times \bar{\psi}), \quad (3)$$

where

$$G_-(s) = i^{-n} \pi^{-n(1/2-s)} \prod_{j=1}^n \Gamma\left(\frac{2-s-\bar{\lambda}_j}{2}\right) \Gamma\left(\frac{s+1-\lambda_j}{2}\right)^{-1}.$$

2.1 Hyper-Kloosterman Sums

Let q be a prime number. For abbreviation, we define $e(x) = \exp(2\pi i x)$. Let k be a positive integer. The $(k-1)$ -dimensional hyper-Kloosterman sum is defined as

$$Kl_k(m, q) = \sum_{\substack{x_1, \dots, x_{k-1} \pmod q \\ (x_1, q) = \dots = (x_{k-1}, q) = 1}} e\left(\frac{x_1 + \dots + x_{k-1} + m \overline{x_1 \dots x_{k-1}}}{q}\right)$$

and we shall note that if $q|m$,

$$Kl_k(m, q) = (-1)^{k-1}. \quad (4)$$

In the degenerate case of $k = 1$, we have the additive character

$$Kl_1(m, q) = e\left(\frac{m}{q}\right).$$

When $k = 2$, the 1-dimensional hyper-Kloosterman sum is identical to the classical Kloosterman sum.

We define $\sum_{\substack{\psi \bmod q \\ \psi(-1)=1}}^*$ as the summation over nontrivial Dirichlet characters ψ of mod q . It is known

$$\sum_{\substack{\psi \bmod q \\ \psi(-1)=1}}^* \tau(\psi)^k \bar{\psi}(m) = \begin{cases} \frac{q-1}{2} (Kl_k(m, q) + Kl_k(-m, q)) - (-1)^k, & \text{if } (m, q) = 1 \\ 0, & \text{otherwise.} \end{cases} \quad (5)$$

It is also known

$$\sum_{\substack{\psi \bmod q \\ \psi(-1)=-1}}^* \tau(\psi)^k \bar{\psi}(m) = \frac{q-1}{2} (Kl_k(m, q) - Kl_k(-m, q)). \quad (6)$$

We will use the family of (multiplicative) Dirichlet characters to recover the additive characters and the hyper-Kloosterman sums by the two formulae above.

3 Even Part

Let q be a prime number, a and $\bar{a}$ two integers with $a\bar{a} \equiv 1 \pmod{q}$. Let k be an integer and $1 \leq k \leq n-1$.

Define two Dirichlet series

$$L_q(s, \pi) = \sum_{q|m, m>0} \frac{A(1, \dots, 1, m)}{m^s}$$

and

$$L_q(s, \tilde{\pi}) = \sum_{q|m, m>0} \frac{A(m, 1, \dots, 1)}{m^s}.$$

Define two Dirichlet series

$$Z(s, q) = \sum_{\substack{\psi \bmod q \\ \psi(-1)=1}}^* \tau(\bar{\psi})^k \psi(a) L(s, \pi \times \psi) + (-1)^k (L(s, \pi) - q L_q(s, \pi))$$

and

$$\tilde{Z}(s, q) = \sum_{\substack{\psi \bmod q \\ \psi(-1)=1}}^* \tau(\psi)^{n-k} \psi(a) L(s, \tilde{\pi} \times \bar{\psi}) + (-1)^{n-k} (L(s, \tilde{\pi}) - q L_q(s, \tilde{\pi})).$$

Lemma 3.1. *We have the identities*

$$Z(s, q) = \frac{1}{2} \sum_{m=1}^{\infty} (Kl_k(am, q) + Kl_k(-am, q)) \frac{A(1, \dots, 1, m)}{m^s}$$

and

$$\tilde{Z}(s, q) = \frac{1}{2} \sum_{m=1}^{\infty} (Kl_{n-k}(\bar{a}m, q) + Kl_{n-k}(-\bar{a}m, q)) \frac{A(m, 1, \dots, 1)}{m^s}.$$

Proof. This follows from (4) and (5), □

For $1 \leq l \leq n-1$, define a Dirichlet polynomial

$$H_l(s, q) = \sum_{i=l}^{n-1} (-1)^i \frac{A(1, \dots, 1, \overbrace{q, 1, \dots, 1}^{\text{position } i})}{q^{is}} + \frac{(-1)^n}{q^{ns}},$$

where q appears on the i^{th} position from the right. We define another Dirichlet polynomial

$$\tilde{H}_l(s, q) = \sum_{i=l}^{n-1} (-1)^i \frac{A(\overbrace{1, \dots, 1, q, 1, \dots, 1}^{\text{position } i})}{q^{is}} + \frac{(-1)^n}{q^{ns}},$$

where q appears on the i^{th} position from the left.

Lemma 3.2. *We have the identities*

$$L(s, \pi) H_l(s, q) = (-1)^l \sum_{m=1}^{\infty} \frac{A(1, \dots, 1, \overbrace{q, 1, \dots, 1, m}^{\text{position } l})}{q^{ls} m^s} \quad (7)$$

for $2 \leq l \leq n-1$ and

$$L(s, \pi) H_1(s, q) = -L_q(s, \pi). \quad (8)$$

Proof. By the Hecke relation

$$\begin{aligned} & A(1, \dots, 1, \overbrace{q, 1, \dots, 1}^{\text{position } i}) A(1, \dots, 1, m) \\ &= \begin{cases} A(1, \dots, 1, \overbrace{q, 1, \dots, 1, m}^{\text{position } i}) + A(1, \dots, 1, \overbrace{q, 1, \dots, 1, m/q}^{\text{position } i+1}), & \text{if } q|m \\ A(1, \dots, 1, \overbrace{q, 1, \dots, 1, m}^{\text{position } i}), & \text{otherwise,} \end{cases} \end{aligned}$$

we have (7) for $2 \leq l \leq n-2$ after multiplying the Dirichlet series $L(s, \pi) H_l(s)$ term by term. Similarly, by the Hecke relation

$$A(1, \dots, 1, q) A(1, \dots, 1, m) = \begin{cases} A(1, \dots, 1, mq) + A(1, \dots, 1, q, m/q), & \text{if } q|m \\ A(1, \dots, 1, mq), & \text{otherwise,} \end{cases}$$

and

$$A(q, 1, \dots, 1) A(1, \dots, 1, m) = \begin{cases} A(q, 1, \dots, 1, m) + A(1, \dots, 1, m/q), & \text{if } q|m \\ A(q, 1, \dots, 1, m), & \text{otherwise,} \end{cases}$$

we have (8), and (7) for $l = n-1$, respectively. □

By a similar proof, we have the following lemma for $\tilde{\pi}$.

Lemma 3.3. *We have the identities*

$$L(s, \tilde{\pi}) \tilde{H}_l(s, q) = (-1)^l \sum_{m=1}^{\infty} \frac{A(\overbrace{m, 1, \dots, 1, q}^{\text{position } l}, 1, \dots, 1)}{q^{ls} m^s}$$

for $2 \leq l \leq n-1$ and

$$L(s, \tilde{\pi}) \tilde{H}_1(s, q) = -L_q(s, \tilde{\pi}).$$

Lemma 3.4. *We have the following identity between Dirichlet polynomials*

$$\begin{aligned} \frac{1}{q} + H_1(s, q) + \sum_{2 \leq l \leq k} (q^{l-1} - q^{l-2}) H_l(s, q) \\ = (-1)^n q^{k-ns} \left(\frac{1}{q} + \tilde{H}_1(1-s, q) + \sum_{2 \leq l \leq n-k} (q^{l-1} - q^{l-2}) \tilde{H}_l(1-s, q) \right). \end{aligned}$$

Proof. The left side equals

$$\frac{1}{q} + \sum_{i=1}^{n-1} (-1)^i q^{\min\{i, k\}-1} \frac{A(1, \dots, 1, \overbrace{q, 1, \dots, 1}^{\text{position } i})}{q^{is}} + \frac{(-1)^n q^{k-1}}{q^{ns}}.$$

The right side equals

$$\begin{aligned} \frac{1}{q} + \sum_{i=1}^{n-1} (-1)^i q^{\min\{i, n-k\}-1} \frac{A(\overbrace{1, \dots, 1, q, 1, \dots, 1}^{\text{position } i})}{q^{i-is}} + \frac{(-1)^n q^{n-k-1}}{q^{n-ns}} \\ = \frac{1}{q} + \sum_{i=1}^{n-1} (-1)^{n-i} q^{\min\{n-i, n-k\}-1} \frac{A(1, \dots, 1, \overbrace{q, 1, \dots, 1}^{\text{position } i})}{q^{(n-i)-(n-i)s}} + \frac{(-1)^n q^{n-k-1}}{q^{n-ns}} \\ = \frac{(-1)^n q^{k-1}}{q^{ns}} + \sum_{i=1}^{n-1} (-1)^{n-i} q^{\min\{0, i-k\}-1} \frac{A(1, \dots, 1, \overbrace{q, 1, \dots, 1}^{\text{position } i})}{q^{-(n-i)s}} + \frac{1}{q} \end{aligned}$$

After multiplying $(-1)^n q^{k-ns}$, we match term by term with the left side. □

3.1 Theorem 3.5

Let $\omega(x) \in C_c^\infty(0, \infty)$ be a test function. Let

$$\tilde{\omega}(s) = \int_0^\infty \omega(x) x^{s-1} dx$$

be the Mellin transform of ω . Define

$$\Omega_+(x) = \frac{1}{2\pi i} \int_{\Re s = -\sigma} \tilde{\omega}(s) G_+(s) x^s ds. \quad (9)$$

We have the Mellin inversion theorem

$$\omega(x) = \frac{1}{2\pi i} \int_{\Re s = \sigma} x^{-s} \tilde{\omega}(s) \, ds. \quad (10)$$

Theorem 3.5. *Let π be an even Maass cusp form for $SL(n, \mathbb{Z})$ with $n \geq 2$. Let $A(m_1, \dots, m_{n-1})$ be the Fourier coefficient of π . Let $\omega \in C_c^\infty(0, \infty)$ be a test function and let q be a prime number, a and $\bar{a}$ two integers with $a\bar{a} \equiv 1 \pmod{q}$. For $1 \leq k \leq n-1$, we have*

$$\begin{aligned} & \frac{1}{2} \sum_{m=1}^{\infty} A(1, \dots, 1, m) (Kl_k(am, q) + Kl_k(-am, q)) \omega(m) \\ & + \sum_{2 \leq l \leq k} \sum_{m=1}^{\infty} (-1)^{l+k} q^{l-1} A(1, \dots, 1, \overbrace{q, 1, \dots, 1}^{\text{position } l}, m) \omega(mq^l) \\ & = \frac{q^k}{2} \sum_{m=1}^{\infty} \frac{A(m, 1, \dots, 1)}{m} (Kl_{n-k}(\bar{a}m, q) + Kl_{n-k}(-\bar{a}m, q)) \Omega_+ \left(\frac{m}{q^n} \right) \\ & + \sum_{2 \leq l \leq n-k} \sum_{m=1}^{\infty} (-1)^{n-k+l} q^{k-1} \frac{A(\overbrace{m, 1, \dots, 1, q}^{\text{position } l}, 1, \dots, 1)}{m} \Omega_+ \left(\frac{m}{q^{n-l}} \right), \end{aligned}$$

where Ω_+ is defined in (9).

Proof. By the functional equations (1) and (2) and Lemma 3.4, we have

$$\begin{aligned} & Z(s, q) + (-1)^k q \sum_{2 \leq l \leq k} (q^{l-1} - q^{l-2}) H_l(s, q) L(s, \pi) \\ & = \sum_{\substack{\psi \pmod{q} \\ \psi(-1)=1}}^* \tau(\bar{\psi})^k \psi(a) L(s, \pi \times \psi) \\ & + (-1)^k \left(L(s, \pi) - q L_q(s, \pi) + q \sum_{2 \leq l \leq k} (q^{l-1} - q^{l-2}) H_l(s, q) L(s, \pi) \right) \\ & = \sum_{\substack{\psi \pmod{q} \\ \psi(-1)=1}}^* \tau(\bar{\psi})^k \psi(a) L(s, \pi \times \psi) \\ & + (-1)^k q \left(\frac{1}{q} + H_1(s, q) + \sum_{2 \leq l \leq k} (q^{l-1} - q^{l-2}) H_l(s, q) \right) L(s, \pi) \\ & = \sum_{\substack{\psi \pmod{q} \\ \psi(-1)=1}}^* \tau(\psi)^{n-k} \psi(a) q^{k-ns} G_+(s) L(1-s, \tilde{\pi} \times \bar{\psi}) \\ & + (-1)^{n-k} q \left(\frac{1}{q} + \tilde{H}_1(1-s, q) + \sum_{2 \leq l \leq n-k} (q^{l-1} - q^{l-2}) \tilde{H}_l(1-s, q) \right) q^{k-ns} G_+(s) L(1-s, \tilde{\pi}) \\ & = q^{k-ns} G_+(s) \left(\tilde{Z}(1-s, q) + (-1)^{n-k} q \sum_{2 \leq l \leq n-k} (q^{l-1} - q^{l-2}) \tilde{H}_l(1-s, q) L(1-s, \tilde{\pi}) \right). \end{aligned}$$

Integrating on both ends with $\tilde{\omega}$ and shifting the integral on the left from $\Re s = -\sigma$ to $\Re s = \sigma$, we have

$$\begin{aligned} & \int_{\Re s = \sigma} \left(Z(s, q) + (-1)^k q \sum_{2 \leq l \leq k} (q^{l-1} - q^{l-2}) H_l(s, q) L(s, \pi) \right) \tilde{\omega}(s) \, ds \\ &= \int_{\Re s = -\sigma} q^{k-ns} G_+(s) \times \\ & \quad \left(\tilde{Z}(1-s, q) + (-1)^{n-k} q \sum_{2 \leq l \leq n-k} (q^{l-1} - q^{l-2}) \tilde{H}_l(1-s, q) L(1-s, \tilde{\pi}) \right) \tilde{\omega}(s) \, ds. \end{aligned}$$

Recalling Lemma 3.1, Lemma 3.2, and Lemma 3.3, we complete the proof with the Mellin inversion (10). $\square$

4 Odd Part

Define

$$Y(s, q) = \sum_{\substack{\psi \pmod q \\ \psi(-1)=-1}}^* \tau(\bar{\psi})^k \psi(a) L(s, \pi \times \psi)$$

and

$$\tilde{Y}(s, q) = \sum_{\substack{\psi \pmod q \\ \psi(-1)=-1}}^* \tau(\bar{\psi})^{n-k} \psi(a) L(s, \tilde{\pi} \times \bar{\psi}).$$

By the function equation (3), we have

$$Y(s, q) = q^{k-ns} G_-(s) \tilde{Y}(1-s, q). \quad (11)$$

Lemma 4.1. *We have the identities*

$$Y(s, q) = \frac{1}{2} \sum_{m=1}^{\infty} (Kl_k(am, q) - Kl_k(-am, q)) \frac{A(1, \dots, 1, m)}{m^s}$$

and

$$\tilde{Y}(s, q) = \frac{1}{2} \sum_{m=1}^{\infty} (Kl_{n-k}(\bar{a}m, q) - Kl_{n-k}(-\bar{a}m, q)) \frac{A(m, 1, \dots, 1)}{m^s}.$$

Proof. This follows from (3). $\square$

Let $\omega(x) \in C_c^\infty(0, \infty)$ be a test function. Let $\tilde{\omega}(s) = \int_0^\infty \omega(x) x^s \, dx / x$ be its Mellin transform. Define

$$\Omega_-(x) = \frac{1}{2\pi i} \int_{\Re s = -\sigma} \tilde{\omega}(s) G_-(s) x^s \, ds. \quad (12)$$

Integrating with $\tilde{\omega}$ on both sides of (11) and shifting the integral on the left to $\Re s = \sigma$, we get

$$\frac{1}{2\pi i} \int_{\Re s = \sigma} Y(s, q) \tilde{\omega}(s) \, ds = \frac{1}{2\pi i} \int_{\Re s = -\sigma} q^{k-ns} G_-(s) \tilde{Y}(1-s, q) \tilde{\omega}(s) \, ds,$$

which, along with Lemma 4.1 and (10), gives us the following theorem.

Theorem 4.2. Let π be an even Maass cusp form for $SL(n, \mathbb{Z})$ with $n \geq 2$. Let $A(m_1, \dots, m_{n-1})$ be the Fourier coefficient of π . Let $\omega \in C_c^\infty(0, \infty)$ be a test function and let q be a prime number, a and $\bar{a}$ two integers with $a\bar{a} \equiv 1 \pmod{q}$. For $1 \leq k \leq n-1$, we have

$$\begin{aligned} \frac{1}{2} \sum_{m=1}^{\infty} A(1, \dots, 1, m) (Kl_k(am, q) - Kl_k(-am, q)) \omega(m) \\ = \frac{q^k}{2} \sum_{m=1}^{\infty} \frac{A(m, 1, \dots, 1)}{m} (Kl_{n-k}(\bar{a}m, q) - Kl_{n-k}(-\bar{a}m, q)) \Omega_{-} \left(\frac{m}{q^n} \right), \end{aligned}$$

where Ω_{-} is defined in (12).

References

- [BuKh] Buttcane, Jack, and Rizwanur Khan. " L^4 -norms of Hecke newforms of large level." *arXiv preprint arXiv:1305.1850* (2013). to appear in *Math. Ann.*, DOI: 10.1007/s00208-014-1142-3.
- [CoPS] Cogdell, James W., and Ilya I. Piatetski-Shapiro. "Converse theorems, functoriality, and applications to number theory." *Proceedings of the International Congress of Mathematicians*, Vol. II (Beijing, 2002), Higher Ed. Press, Beijing, 2002: 119-128.
- [Go] Goldfeld, Dorian. *Automorphic forms and L-functions for the group $GL(n, \mathbb{R})$* . Cambridge University Press, 2006.
- [GoLi1] Goldfeld, Dorian, and Xiaoqing Li. "Voronoi formulas on $GL(n)$." *International Mathematics Research Notices* 2006 (2006): 86295.
- [GoLi2] Goldfeld, Dorian, and Xiaoqing Li. "The Voronoi formula for $GL(n, \mathbb{R})$." *International Mathematics Research Notices* 2008 (2008): rnm144.
- [IcTe] Ichino, Atsushi, and Nicolas Templier. "On the Voronoi formula for $GL(n)$." *American Journal of Mathematics* 135, no. 1 (2013): 65-101.
- [Li] Li, Xiaoqing. "Bounds for $GL(3) \times GL(2)$ L-functions and $GL(3)$ L-functions." *Annals of Mathematics* 173 (2011): 301-336.
- [Mi] Miller, Stephen David. "Cancellation in additively twisted sums on $GL(n)$." *American journal of mathematics* 128, no. 3 (2006): 699-729.
- [MiSc1] Miller, Stephen D., and Wilfried Schmid. "Automorphic distributions, L-functions, and Voronoi summation for $GL(3)$." *Annals of mathematics* (2006): 423-488.
- [MiSc2] Miller, Stephen D., and Wilfried Schmid. "A general Voronoi summation formula for $GL(n, \mathbb{Z})$." In *Geometry and analysis*. No. 2, volume 18 of Adv. Lect. Math. (ALM), pages 173-224. Int. Press, Somerville, MA, 2011.

FAN ZHOU
Department of Mathematics
The Ohio State University
Columbus, OH 43210, USA
zhou.1406@math.osu.edu